

1. In ΔOPQ , right-angled at P, $OP = 7 \text{ cm}$ and $OQ - PQ = 1 \text{ cm}$, then the values of $\sin Q$.
 (a) $\frac{7}{25}$ (b) $\frac{24}{25}$ (c) 1 (d) none of the these

2. If $\sin A = \frac{24}{25}$, then the value of $\cos A$ is
 (a) $\frac{7}{25}$ (b) $\frac{24}{25}$ (c) 1 (d) none of the these

3. In ΔABC , right-angled at B, $AB = 5 \text{ cm}$ and $\angle ACB = 30^\circ$ then the length of the side BC is
 (a) $5\sqrt{3}$ (b) $2\sqrt{3}$ (c) 10 cm (d) none of these

4. In ΔABC , right-angled at B, $AB = 5 \text{ cm}$ and $\angle ACB = 30^\circ$ then the length of the side AC is
 (a) $5\sqrt{3}$ (b) $2\sqrt{3}$ (c) 10 cm (d) none of these

5. The value of $\frac{2 \tan 30^\circ}{1 + \tan^2 30^\circ}$ is
 (a) $\sin 60^\circ$ (b) $\cos 60^\circ$ (c) $\tan 60^\circ$ (d) $\sin 30^\circ$

6. The value of $\frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ}$ is
 (a) $\tan 90^\circ$ (b) 1 (c) $\sin 45^\circ$ (d) 0

7. $\sin 2A = 2 \sin A$ is true when $A =$
 (a) 0° (b) 30° (c) 45° (d) 60°

8. The value of $\frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ}$ is
 (a) $\sin 60^\circ$ (b) $\cos 60^\circ$ (c) $\tan 60^\circ$ (d) $\sin 30^\circ$

9. $9 \sec^2 A - 9 \tan^2 A =$
 (a) 1 (b) 9 (c) 8 (d) 0

10. $(1 + \tan A + \sec A)(1 + \cot A - \operatorname{cosec} A) =$
 (a) 0 (b) 1 (c) 2 (d) -1

11. $(\sec A + \tan A)(1 - \sin A) =$
 (a) $\sec A$ (b) $\sin A$ (c) $\operatorname{cosec} A$ (d) $\cos A$

12. $\frac{1 + \tan^2 A}{1 + \cot^2 A} =$
 (a) $\sec^2 A$ (b) -1 (c) $\cot^2 A$ (d) $\tan^2 A$

1. If $\sin 3A = \cos(A - 26^\circ)$, where $3A$ is an acute angle, find the value of A .
 (a) 29° (b) 30° (c) 26° (d) 36°

2. If $\tan 2A = \cot(A - 18^\circ)$, where $2A$ is an acute angle, find the value of A .
 (a) 29° (b) 30° (c) 26° (d) none of these

3. If $\sec 4A = \operatorname{cosec}(A - 20^\circ)$, where $4A$ is an acute angle, find the value of A .
 (a) 22° (b) 25° (c) 26° (d) none of these

4. The value of $\tan 48^\circ \tan 23^\circ \tan 42^\circ \tan 67^\circ$ is
 (a) 1 (b) 9 (c) 8 (d) 0

5. If ΔABC is right angled at C, then the value of $\cos(A + B)$ is
 (a) 0 (b) 1 (c) $\frac{1}{2}$ (d) n.d.

6. The value of the expression $\left[\frac{\sin^2 22^\circ + \sin^2 68^\circ}{\cos^2 22^\circ + \cos^2 68^\circ} + \sin^2 63^\circ + \cos 63^\circ \sin 27^\circ \right]$ is
 (a) 3 (b) 0 (c) 1 (d) 2

7. If $\cos A = \frac{24}{25}$, then the value of $\sin A$ is
 (a) $\frac{7}{25}$ (b) $\frac{24}{25}$ (c) 1 (d) none of the these

8. If ΔABC is right angled at B, then the value of $\cos(A + C)$ is
 (a) 0 (b) 1 (c) $\frac{1}{2}$ (d) n.d.

9. If $\tan A = \frac{4}{3}$, then the value of $\cos A$ is
 (a) $\frac{3}{5}$ (b) $\frac{4}{3}$ (c) 1 (d) none of the these

10. If ΔABC is right angled at C, in which $AB = 29$ units, $BC = 21$ units and $\angle ABC = \alpha$. Determine the values of $\cos^2 \alpha + \sin^2 \alpha$ is
 (a) 0 (b) 1 (c) $\frac{1}{2}$ (d) n.d.

11. In a right triangle ABC, right-angled at B, if $\tan A = 1$, then the value of $2 \sin A \cos A$ =
 (a) 0 (b) 1 (c) $\frac{1}{2}$ (d) n.d.

12. Given $15 \cot A = 8$, then $\sin A$ =
 (a) $\frac{3}{5}$ (b) $\frac{4}{3}$ (c) 1 (d) none of the these

1. In a triangle PQR, right-angled at Q, $PR + QR = 25$ cm and $PQ = 5$ cm, then the value of $\sin P$ is
 (a) $\frac{7}{25}$ (b) $\frac{24}{25}$ (c) 1 (d) none of the these
2. In a triangle PQR, right-angled at Q, $PQ = 3$ cm and $PR = 6$ cm, then $\angle QPR =$
 (a) 0° (b) 30° (c) 45° (d) 60°
3. If $\sin(A - B) = \frac{1}{2}$ and $\cos(A + B) = \frac{1}{2}$, then the value of A and B, respectively are
 (a) 45° and 15° (b) 30° and 15° (c) 45° and 30° (d) none of these
4. If $\sin(A - B) = 1$ and $\cos(A + B) = 1$, then the value of A and B, respectively are
 (a) 45° and 15° (b) 30° and 15° (c) 45° and 30° (d) none of these
5. If $\tan(A - B) = \frac{1}{\sqrt{3}}$ and $\tan(A + B) = \sqrt{3}$, then the value of A and B, respectively are
 (a) 45° and 15° (b) 30° and 15° (c) 45° and 30° (d) none of these
6. If $\cos(A - B) = \frac{\sqrt{3}}{2}$ and $\sin(A + B) = 1$, then the value of A and B, respectively are
 (a) 45° and 15° (b) 30° and 15° (c) 60° and 30° (d) none of these
7. The value of $2\cos^2 60^\circ + 3\sin^2 45^\circ - 3\sin^2 30^\circ + 2\cos^2 90^\circ$ is
 (a) 1 (b) 5 (c) $5/4$ (d) none of these
8. $\sin 2A = 2 \sin A \cos A$ is true when A =
 (a) 0° (b) 30° (c) 45° (d) any angle
9. $\sin A = \cos A$ is true when A =
 (a) 0° (b) 30° (c) 45° (d) any angle
10. If $\sin A = \frac{1}{2}$, then the value of $3\cos A - 4\cos^3 A$ is
 (a) 0 (b) 1 (c) $1/2$ (d) n.d.
11. If $3\cot A = 4$, then the value of $\cos^2 A - \sin^2 A$ is
 (a) $\frac{3}{4}$ (b) $\frac{7}{25}$ (c) $1/2$ (d) $\frac{24}{25}$
12. If $3\tan A = 4$, then the value of $\frac{3\sin A + 2\cos A}{3\sin A - 2\cos A}$ is
 (a) 1 (b) $\frac{7}{25}$ (c) 3 (d) $\frac{24}{25}$

1. Value of θ , for $\sin 2\theta = 1$, where $0^\circ < \theta < 90^\circ$ is:
 (a) 30° (b) 60° (c) 45° (d) 135° .

2. Value of $\sec^2 26^\circ - \cot^2 64^\circ$ is:
 (a) 1 (b) -1 (c) 0 (d) 2

3. Product $\tan 1^\circ \cdot \tan 2^\circ \cdot \tan 3^\circ \dots \tan 89^\circ$ is:
 (a) 1 (b) -1 (c) 0 (d) 90

4. $\sqrt{1 + \tan^2 \theta}$ is equal to:
 (a) $\cot \theta$ (b) $\cos \theta$ (c) $\operatorname{cosec} \theta$ (d) $\sec \theta$

5. If $A + B = 90^\circ$, $\cot B = \frac{3}{4}$ then $\tan A$ is equal to;
 (a) $\frac{3}{4}$ (b) $\frac{4}{3}$ (c) $\frac{1}{4}$ (d) $\frac{1}{3}$

6. Maximum value of $\frac{1}{\operatorname{cosec} \theta}$, $0^\circ < \theta < 90^\circ$ is:
 (a) 1 (b) -1 (c) 2 (d) $\frac{1}{2}$

7. If $\cos \theta = \frac{1}{2}$, $\sin \phi = \frac{1}{2}$ then value of $\theta + \phi$ is
 (a) 30° (b) 60° (c) 90° (d) 120° .

8. If $\sin(A + B) = 1 = \cos(A - B)$ then
 (a) $A = B = 90^\circ$ (b) $A = B = 0^\circ$ (c) $A = B = 45^\circ$ (d) $A = 2B$

9. The value of $\sin 60^\circ \cos 30^\circ - \cos 60^\circ \sin 30^\circ$ is
 (a) 1 (b) -1 (c) 0 (d) none of these

10. The value of $2\sin^2 30^\circ - 3\cos^2 45^\circ + \tan^2 60^\circ + 3\sin^2 90^\circ$ is
 (a) 1 (b) 5 (c) 0 (d) none of these